

Nonperturbatively Regulating Chiral Gauge Theories

DMG and David B. Kaplan
arXiv:1511.03649

Motivation: Lattice Regulate Chiral Gauge Theory

Big Question 1: Do chiral gauge theories (χ GT) make sense beyond perturbation theory?

- Only known χ GT is the Standard Model Electroweak sector
- What are the requirements to have a well-defined χ GT

Big Question 2: What are the properties of strongly interacting chiral gauge theories?

- High energy extensions of the Standard Model

To answer these, must first find a nonperturbative regulator.

Motivation: Lattice Regulate Chiral Gauge Theory

Vector Theory (QED, QCD)

- **Real** fermion representation
- Gauge symmetries **allow** fermion mass term
- Gauge invariant massive regulator (Pauli-Villars) **can** be used
- **Known** lattice regulator

Motivation: Lattice Regulate Chiral Gauge Theory

Vector Theory (QED, QCD)

- **Real** fermion representation
- Gauge symmetries **allow** fermion mass term
- Gauge invariant massive regulator (Pauli-Villars) **can** be used
- **Known** lattice regulator

Chiral Theory (Electroweak)

- **Complex** fermion representation
- Gauge symmetries **forbid** fermion mass term
- Gauge invariant massive regulator **cannot** be used
- **No widely accepted** lattice regulator (Eichten and Preskill '86, Narayanan and Neuberger '94, Lüscher '99, etc)

Motivation: Lattice Regulate Chiral Gauge Theory

Vector Theory (QED, QCD)

- **Real** fermion representation
- Gauge symmetries **allow** fermion mass term
- Gauge invariant massive regulator (Pauli-Villars) **can** be used
- **Known** lattice regulator

Chiral Theory (Electroweak)

- **Complex** fermion representation
- Gauge symmetries **forbid** fermion mass term
- Gauge invariant massive regulator **cannot** be used
- **No widely accepted** lattice regulator (Eichten and Preskill '86, Narayanan and Neuberger '94, Lüscher '99, etc)

Is this a technical issue or indicative of new physics and thus a problem with the Standard Model?

Technical Question: Define Measure for χ GT

Observables are calculated by integrating over gauge fields with some measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

- $F(A)$ is the observable
- $S(A)$ is gauge action (Maxwell or Yang Mills)
- $\Delta(A)$ is due to fermions

Technical Question: Define Measure for χ GT

Observables are calculated by integrating over gauge fields with some measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

- $F(A)$ is the observable
- $S(A)$ is gauge action (Maxwell or Yang Mills)
- $\Delta(A)$ is due to fermions
 - $\Delta(A)$ for Dirac fermion is well-known

$$\Delta_{DF}(A) = \det \not{D}(A)$$

- But it is not well know how to define $\Delta(A)$ for chiral fermion

$$\Delta_{\chi F} \Delta_{\chi F}^* = \Delta_{DF}$$

Technical Question: Define Measure for χ GT

What is the fermionic contribution to the measure for χ GT?

- Need definition so that effective action is local and analytic


$$\Delta_{DF}(A) = \prod \lambda_i \leftarrow \text{Dirac operator eigenvalues}$$

Technical Question: Define Measure for χ GT

What is the fermionic contribution to the measure for χ GT?

- Need definition so that effective action is local and analytic

$$\Delta_{DF}(A) = \prod \lambda_i \leftarrow \text{Dirac operator eigenvalues}$$




$$\Delta_{\chi F}(A) = \prod_{\lambda_j > 0} \lambda_j$$

Technical Question: Define Measure for χ GT


What is the fermionic contribution to the measure for χ GT?

- Need definition so that effective action is local and analytic

$$\Delta_{DF}(A) = \prod \lambda_i \leftarrow \text{Dirac operator eigenvalues}$$



$$\Delta_{\chi F}(A) = \prod_{\lambda_j > 0} \lambda_j$$



$$\Delta_{\chi F}(A) = \prod_{\lambda_j(A^*) > 0} \lambda_j$$

Motivation: Lattice Regulate Chiral Gauge Theory

Continuum Field Theory

- Theories with chiral symmetries can have anomalies
- Standard Model contains global anomalies
- Chiral gauge theories only well-behaved if no gauge anomalies

Motivation: Lattice Regulate Chiral Gauge Theory

Continuum Field Theory

- Theories with chiral symmetries can have anomalies
- Standard Model contains global anomalies
- Chiral gauge theories only well-behaved if no gauge anomalies

Lattice Field Theory

- No anomalies in system with finite degrees of freedom
- Lattice must explicitly break global chiral symmetry to reproduce anomaly
- Lattice must somehow distinguish anomalous and anomaly-free gauge theories

Motivation: Lattice Regulate Chiral Gauge Theory

Continuum Field Theory

- Theories with chiral symmetries can have anomalies
- Standard Model contains global anomalies
- Chiral gauge theories only well-behaved if no gauge anomalies

Lattice Field Theory

- No anomalies in system with finite degrees of freedom
- Lattice must explicitly break global chiral symmetry to reproduce anomaly
- Lattice must somehow distinguish anomalous and anomaly-free gauge theories

How does one construct a lattice theory that has the correct continuum behavior?

Motivation: Lattice Regulate Chiral Gauge Theory

Choice A: Explicit Gauge Violation

- Lattice theory is not gauge invariant
- Gauge invariance must be restored in continuum limit
- Sensible continuum limit only exists for anomaly-free theories

Motivation: Lattice Regulate Chiral Gauge Theory

Choice A: Explicit Gauge Violation

- Lattice theory is not gauge invariant
- Gauge invariance must be restored in continuum limit
- Sensible continuum limit only exists for anomaly-free theories

Choice B: Maintain Gauge Invariance

- Lattice theory is gauge invariant
- Must have a mechanism to distinguish anomalous and anomaly-free gauge theories

Motivation: Lattice Regulate Chiral Gauge Theory

Choice A: Explicit Gauge Violation

- Lattice theory is not gauge invariant
- Gauge invariance must be restored in continuum limit
- Sensible continuum limit only exists for anomaly-free theories

Choice B: Maintain Gauge Invariance

- Lattice theory is gauge invariant
- Must have a mechanism to distinguish anomalous and anomaly-free gauge theories

We will go with Choice B

For our construction, only anomaly-free theories are local

Steps to Define Measure for χ GT

Basic building block is Dirac fermion, in order to have well-defined eigenvalue problem

1. Global chiral symmetry (massless Dirac fermions)
2. Decouple mirror fermions
3. Mechanism for distinguishing anomalous versus anomaly free fermion representation

Steps to Define Measure for χ GT

Basic building block is Dirac fermion, in order to have well-defined eigenvalue problem


1. Global chiral symmetry (massless Dirac fermions)
2. Decouple mirror fermions
3. Mechanism for distinguishing anomalous versus anomaly free fermion representation

Global Chiral Symmetries

Domain Wall Fermions (DWF)

(Kaplan, '92)

- Introduce extra (compact) dimension, x_5
- Fermion mass depends on x_5
- Massless modes localized on mass defects
- Gauge fields independent of x_5
- Anomaly due to bulk fermions carrying charge between mass defects
- Condensed matter physicists would call this a topological insulator



Global Chiral Symmetries


DWF always give rise to a vector gauge theory

- DWF 5d action is equivalent to action for an infinite number of 4d fermions
- If discretize extra dimension, x_5 is a flavor quantum number

$$\bar{\psi} \gamma_5 \partial_5 \psi \rightarrow \bar{\psi}_n \gamma_5 (\psi_{n+1} - \psi_n)$$



Every flavor must be in same gauge group representation



Steps to Define Fermion Measure for χ GT


Basic building block is Dirac fermion, in order to have well-defined eigenvalue problem

1. Global chiral symmetry (massless Dirac fermions)
2. Decouple mirror fermions
3. Mechanism for distinguishing anomalous versus anomaly free fermion representation

Gauged Chiral Symmetries (Previous Attempt)

Idea: Localize Gauge Fields at one defect

- Waveguide Model
- Gauge fields **depend** on x_5
- Need spontaneous symmetry breaking to preserve gauge invariance
- New RH mode appears at location of SSB
- Spectrum has **Dirac** fermions
Golterman, Jansen, Vink 1993



Gauged Chiral Symmetries (New Attempt)

New Idea: Localize gauge fields around one defect via smearing
(DMG and Kaplan, '15II)

Smeared Gauge Fields (Narayanan and Neuberger, '06; Lüscher, '11, etc)


- Utilizes extra dimension
- Start with any gauge field, A_μ
- Extend gauge field into the bulk via particular flow equation

$$\text{Flow Eq: } \partial_s \bar{A}_\mu = D_\nu \bar{F}_{\nu\mu} \quad \text{BC: } \bar{A}_\mu(x, 0) = A_\mu(x)$$

- Behaves like heat equation
- **Damps out high momentum modes**

Flow Equation: 2d/3d QED Example

4d World



Write A_μ in terms of gauge and physical degree of freedom

$$\bar{A}_\mu = \partial_\mu \bar{\omega} + \epsilon_{\mu\nu} \partial_\nu \bar{\lambda}$$

Flow Eqs.


$$\partial_s \bar{\lambda} = \square \bar{\lambda} \quad \partial_s \bar{\omega} = 0$$

Flow in extra dimension damps out high momenta modes

Combine Domain Wall Fermions and Smearing

New Idea: Localize gauge fields at one defect via smearing

- Gauge field at $s=0$ is quantum gauge field $A_\mu(x)$
- Bulk gauge field $\bar{A}_\mu(x,s)$ obeys flow equation
- Flow is symmetric around $s=0$
- RH modes have soft form factor coupling to physical degrees of freedom




Combine Domain Wall Fermions and Smearing

New Idea: Localize gauge fields at one defect via smearing

- Gauge field at $s=0$ is quantum gauge field $A_\mu(x)$
- Bulk gauge field $\bar{A}_\mu(x,s)$ obeys flow equation
- Flow is symmetric around $s=0$
- RH modes have soft form factor coupling to physical degrees of freedom

$$e^{-\xi p^2 L/\Lambda}$$



Combine Domain Wall Fermions and Smearing


New Idea: Localize gauge fields at one defect via smearing

- Gauge field at $s=0$ is quantum gauge field $A_\mu(x)$
- Bulk gauge field $\bar{A}_\mu(x,s)$ obeys flow equation
- Flow is symmetric around $s=0$
- RH modes have soft form factor coupling to physical degrees of freedom

$$e^{-\xi p^2 L/\Lambda}$$

Photon Momentum

Wall separation




Combine Domain Wall Fermions and Smearing

New Idea: Localize gauge fields at one defect via smearing

- Gauge field at $s=0$ is quantum gauge field $A_\mu(x)$
- Bulk gauge field $\bar{A}_\mu(x,s)$ obeys flow equation
- Flow is symmetric around $s=0$
- RH modes have soft form factor coupling to physical degrees of freedom

$$e^{-\xi p^2 L / \Lambda}$$

Flow parameter Bulk fermion mass




Combine Domain Wall Fermions and Smearing

New Idea: Localize gauge fields at one defect via smearing

- Gauge field at $s=0$ is quantum gauge field $A_\mu(x)$
- Bulk gauge field $\bar{A}_\mu(x,s)$ obeys flow equation
- Flow is symmetric around $s=0$
- RH modes have soft form factor coupling to physical degrees of freedom

$$e^{-\xi p^2 L/\Lambda}$$

- LH and RH modes couple equally to gauge degrees of freedom




Steps to Define Fermion Measure for χ GT

Basic building block is Dirac fermion, in order to have well-defined eigenvalue problem

1. Global chiral symmetry (massless Dirac fermions)
2. Decouple mirror fermions
3. Mechanism for distinguishing anomalous versus anomaly free fermion representation


Anomalies and Callan-Harvey Mechanism (Callan and Harvey, '84)

Integrating out bulk fermions generates a Chern-Simons term



Anomalies and Callan-Harvey Mechanism (Callan and Harvey, '84)

Integrating out bulk fermions generates a Chern-Simons term



Anomalies and Callan-Harvey Mechanism (Callan and Harvey, 1984)

- Integrating out bulk fermions generates a Chern-Simons term
- In 3 dimensions, the Chern Simons action is

$$S_3^{\text{bulk}} = c_3 \frac{\Lambda}{|\Lambda|} \int (\epsilon(s) - 1) \text{Tr} \left(\bar{F} \bar{A} - \frac{1}{3} \bar{A}^3 \right)$$

Anomalies and Callan-Harvey Mechanism (Callan and Harvey, 1984)

- Integrating out bulk fermions generates a Chern-Simons term
- In 3 dimensions, the Chern Simons action is

$$S_3^{\text{bulk}} = c_3 \frac{\Lambda}{|\Lambda|} \int (\epsilon(s) - 1) \text{Tr} \left(\bar{F} \bar{A} - \frac{1}{3} \bar{A}^3 \right)$$

CS only depends on sign
of domain wall mass

Fermion Contribution

Pauli Villars Contribution

Anomalies and Callan-Harvey Mechanism (Callan and Harvey, 1984)

- Integrating out bulk fermions generates a Chern-Simons term
- In 3 dimensions, the Chern Simons action is

$$S_3^{\text{bulk}} = c_3 \frac{\Lambda}{|\Lambda|} \int (\epsilon(s) - 1) \text{Tr} \left(\bar{F} \bar{A} - \frac{1}{3} \bar{A}^3 \right)$$

CS only depends on sign
of domain wall mass

Fermion Contribution

Pauli Villars Contribution

- This approximation is only valid far away from domain wall

Anomalies and Callan-Harvey Mechanism

- Consider 3 dimensional QED with flowed gauge fields

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- No gauge field in 3rd dimension

Anomalies and Callan-Harvey Mechanism

- Consider 3 dimensional QED with flowed gauge fields

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- No gauge field in 3rd dimension
- Effective two point function is nonlocal

$$\Gamma(r) = \left(\delta^2(r) - \frac{\mu^2}{4\pi} e^{-\mu^2 r^2/4} \right) \quad \mu \equiv \sqrt{\frac{\Lambda}{\xi L}}$$

- When flow is turned off ($\mu \rightarrow \infty$), Γ vanishes

Anomalies and Callan-Harvey Mechanism

- Consider 3 dimensional QED with flowed gauge fields

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- No gauge field in 3rd dimension
- Effective two point function is nonlocal

$$\Gamma(r) = \left(\delta^2(r) - \frac{\mu^2}{4\pi} e^{-\mu^2 r^2 / 4} \right)$$

$$\mu \equiv \sqrt{\frac{\Lambda}{\xi L}}$$

Serves as an IR cutoff

- When flow is turned off ($\mu \rightarrow \infty$), Γ vanishes

Anomaly Cancellation and Nonlocality

- DWF with flowed gauge fields gives rise to a nonlocal 2d theory

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- If include multiple fields, Chern Simons prefactor is

$$\sum_i e_i^2 \frac{\Lambda_i}{|\Lambda_i|}$$

- The theory is local if this prefactor vanishes

Anomaly Cancellation and Nonlocality

- DWF with flowed gauge fields gives rise to a nonlocal 2d theory

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- If include multiple fields, Chern Simons prefactor is

$$\sum_i e_i^2 \frac{\Lambda_i}{|\Lambda_i|}$$

Fermion Chirality

- The theory is local if this prefactor vanishes

Anomaly Cancellation and Nonlocality

- DWF with flowed gauge fields gives rise to a nonlocal 2d theory

$$S_3^{\text{bulk}} = 2e^2 c_3 \frac{\Lambda}{|\Lambda|} \int dx^2 dy^2 \left(\frac{\partial_\mu \partial_\alpha}{\square} A_\alpha(x) \right) \Gamma(x-y) \left(\frac{\partial_\mu \partial_\beta}{\square} \epsilon_{\beta\gamma} A_\gamma(y) \right)$$

- If include multiple fields, Chern Simons prefactor is

$$\sum_i e_i^2 \frac{\Lambda_i}{|\Lambda_i|}$$

Fermion Chirality

- The theory is local if this prefactor vanishes

This is exactly equivalent to the requirement that the chiral fermions be in an anomaly free representation

Steps to Define Fermion Measure for χ GT

Basic building block is Dirac fermion, in order to have well-defined eigenvalue problem

1. Global chiral symmetry (massless Dirac fermions)
 - Domain Wall Fermions
2. Decouple mirror fermions
 - Smeared Gauge Fields
3. Mechanism for distinguishing anomalous versus anomaly free fermion representation
 - Theory is local if fermions are in an anomaly free representation

Proposal for Chiral Gauge Theory Measure

Recall that the goal is to be able to define a chiral fermion measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

Our proposal:

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]}$$

$$\partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

Proposal for Chiral Gauge Theory Measure

Recall that the goal is to be able to define a chiral fermion measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

Our proposal:

$$\Delta(A) = \prod_i \frac{\overbrace{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}^{\text{DWF}}}{\underbrace{\det [\not{D}(\bar{A}) - \Lambda_i]}_{\text{Pauli-Villars}}}$$

$$\partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

Proposal for Chiral Gauge Theory Measure

Recall that the goal is to be able to define a chiral fermion measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

Our proposal:

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]}$$

One factor for each
species of fermion

5d Dirac operator
with flowed gauge field

$$\partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

Proposal for Chiral Gauge Theory Measure

Recall that the goal is to be able to define a chiral fermion measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

Our proposal:

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]}$$

$$\partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

Proposal for Chiral Gauge Theory Measure

Recall that the goal is to be able to define a chiral fermion measure

$$\langle F(A) \rangle = \frac{\int [DA] e^{-S(A)} \Delta(A) F(A)}{\int [DA] e^{-S(A)} \Delta(A)}$$

Our proposal:

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]}$$

$$\partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

- Mirror fermions decouple for large wall separation
- Target d-dimensional theory is local if fermions are in an anomaly free representation
- Effective action is what one would expect for chiral fermion (did not show here)

Open Questions

Open Question 1: What is the behavior of topological gauge configurations

- Do the mirrors decouple from topological gauge configurations?
- Can the Standard Model particles exchange energy and momenta with the mirror fermions

Open Questions

Open Question 1: What is the behavior of topological gauge configurations

- Do the mirrors decouple from topological gauge configurations?
- Can the Standard Model particles exchange energy and momenta with the mirror fermions

Open Question 2: Does the theory have a well defined Hamiltonian and Minkowski space path integral

- Is the Hamiltonian bounded and hermitian?
- Is the S matrix unitary?
- Is the theory casual?

Open Questions


Open Question 1: What is the behavior of topological gauge configurations

- Do the mirrors decouple from topological gauge configurations?
- Can the Standard Model particles exchange energy and momenta with the mirror fermions


Open Question 2: Does the theory have a well defined Hamiltonian and Minkowski space path integral

- Is the Hamiltonian bounded and hermitian?
- Is the S matrix unitary?
- Is the theory casual?

Topological Gauge Configurations - Weak Coupling




Topological Gauge Configurations - Weak Coupling




At weak coupling, instanton contribution is most important

- Flow does not affect location of instantons
- Correlation between location of instantons on the two boundaries allows for exchange of energy/momentum
- Highly suppressed process, so difficult to observe

Topological Gauge Configurations - Strong Coupling



Topological Gauge Configurations - Strong Coupling



At strong coupling, need to include instanton-anti instanton pairs

- I-A pairs DO NOT satisfy equations of motion
- If flow for sufficiently long time, all pairs will annihilate
- If no correlation between location of instantons on the two boundary, standard fermions and Fluff do not exchange energy/momentum

Summary

- Proposal for fermion measure for chiral gauge theory

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]} \quad \partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

- Combines domain wall fermions and gauge field smearing
- Local theory if chiral fermion representation is anomaly free
- Mirror fermions decouple due to exponentially soft form factors to gauge fields

Summary

- Proposal for fermion measure for chiral gauge theory

$$\Delta(A) = \prod_i \frac{\det [\not{D}(\bar{A}) - \Lambda_i \epsilon(s)]}{\det [\not{D}(\bar{A}) - \Lambda_i]} \quad \partial_s \bar{A}_\mu = \frac{\xi \epsilon(s)}{|\Lambda|} D_\nu \bar{F}_{\nu\mu}$$

- Combines domain wall fermions and gauge field smearing
 - Local theory if chiral fermion representation is anomaly free
 - Mirror fermions decouple due to exponentially soft form factors to gauge fields
-
- Important open questions remain about this proposal
 - Is there Fluff hiding in the Standard Model?